

EC analytical columns NUCLEODUR^{®} 100-5 C₁₈ ec, 5 µm

Octadecyl phases, 17.5 % C, particle size 5 µm.
 Eluent in column: Acetonitrile/water
 Particle type: Fully porous
 Endcapped: Yes
 Carbon content: 17.5 %
 pH stability: 1 ... 9
 Max. temp.resistance: 60 °C

| Method | Material column | USP listing | Separation system | Particle size µm | Int. Ø mm | Length mm | Volume column ml | PK | Cat. No. |
|------------|-----------------|-------------|-------------------|---------------------|--------------|--------------|---------------------|----|-----------|
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 2 | 50 | 0.15 | 1 | 4.004 368 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 3 | 50 | 0.35 | 1 | 4.004 369 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 4 | 50 | 0.62 | 1 | 4.004 370 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 4.6 | 50 | 0.83 | 1 | 4.004 371 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 4.6 | 100 | 1.65 | 1 | 4.006 934 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 2 | 125 | 0.37 | 1 | 9.003 816 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 3 | 125 | 0.87 | 1 | 9.003 817 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 4 | 125 | 1.56 | 1 | 9.003 818 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 4.6 | 125 | 2.06 | 1 | 9.003 819 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 4.6 | 150 | 2.5 | 1 | 9.003 820 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 2 | 250 | 0.8 | 1 | 9.003 821 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 3 | 250 | 1.75 | 1 | 9.003 822 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 4 | 250 | 3.15 | 1 | 9.003 823 |
| Analytical | Stainless steel | L1 | Reverse phase | 5 | 4.6 | 250 | 4.150 | 1 | 9.003 824 |